

# EVENLY DIVISIBLE RATIONAL APPROXIMATIONS OF QUADRATIC IRRATIONALITIES

DAN CARMON

**ABSTRACT.** In a recent paper of Blomer, Bourgain, Radziwiłł and Rudnick [1], the authors proved the existence of small gaps between eigenvalues of the Laplacian in a rectangular billiard with sides  $\pi$  and  $\pi/\sqrt{\alpha}$ , i.e. numbers of the form  $\alpha m^2 + n^2$ , whenever  $\alpha$  is a quadratic irrationality of certain types. In this note, we extend their results to all positive quadratic irrationalities  $\alpha$ .

## 1. INTRODUCTION

In [1], the authors investigated the minimal gaps between energy levels of the eigenvalues of the Laplacian of a billiard in a rectangle with width  $\pi/\sqrt{\alpha}$  and height  $\pi$ . These eigenvalues are of the form  $\alpha m^2 + n^2$ , where  $m, n \geq 1$  are integers. For positive irrational  $\alpha$ , these levels belong to a simple spectrum  $0 < \lambda_1 < \lambda_2 < \dots$ , with growth  $\lambda_N \sim \frac{4\sqrt{\alpha}}{\pi}N$ . The minimal gap function is then defined as

$$\delta_{\min}^{(\alpha)}(N) = \min(\{\lambda_{i+1} - \lambda_i : 1 \leq i < N\}).$$

The authors prove several lower and upper bounds on the growth rate of  $\delta_{\min}^{(\alpha)}(N)$ , for various families and sets of irrational  $\alpha$ . Amongst these is [1, Theorem 6.1]:

**Theorem 1.1.** *For all positive real quadratic irrationalities of the form*

$$(1.1) \quad \alpha = \alpha(x; a, b, \epsilon, r) = r \cdot \left( \frac{x + \sqrt{x^2 + 4\epsilon}}{2} \right)^a \cdot \left( \sqrt{x^2 + 4\epsilon} \right)^b$$

with

$$a \in \mathbb{Z}, \quad b = 0, 1, \quad x \in \mathbb{Z} \setminus \{0\}, \quad \epsilon = \pm 1, \quad r \in \mathbb{Q}^\times,$$

we have  $\delta_{\min}^{(\alpha)}(N) \ll_{\alpha, \epsilon} N^{-1+\epsilon}$  infinitely often, for any  $\epsilon > 0$ .

Furthermore, the authors remark that when  $b = 0$  and  $a$  is even, they have in fact obtained the stronger result that  $\delta_{\min}^{(\alpha)}(N) \ll_{\alpha} N^{-1}$  for all  $N$ . We refer the reader to [1] for a more detailed introduction to the problem, as well as the motivation for these upper bounds.

---

*Date:* December 6, 2016.

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement n° 320755.

Our goal in this paper is to generalize the result above to any positive quadratic irrationality. Here on, we shall denote by  $K = \mathbb{Q}[\sqrt{D}]$  – a real quadratic number field,  $\mathcal{O}_K$  – its ring of integers, and for any  $\omega \in K$ , we shall denote its conjugate in  $K$  by  $\bar{\omega}$ , and its norm and trace (over  $\mathbb{Q}$ ) by  $N(\omega) = \omega\bar{\omega}$ ,  $\text{tr}(\omega) = \omega + \bar{\omega}$ . In section 3, we prove:

**Theorem 1.2.** *Let  $K = \mathbb{Q}[\sqrt{D}]$  be any real quadratic number field, where  $D \in \mathbb{N}$  is square-free. Let  $0 < \alpha \in K$  be a positive element of  $K$ . Then  $\delta_{\min}^{(\alpha)}(N) \ll_{\alpha, \varepsilon} N^{-1+\varepsilon}$  infinitely often, for any  $\varepsilon > 0$ . Furthermore, if  $\alpha \in \mathbb{Q}_{>0} \cdot (K^\times)^2$ ,<sup>1</sup> then  $\delta_{\min}^{(\alpha)}(N) \ll N^{-1}$  for all  $N$ .*

One of the key lemmas in [1] relates small gaps between the  $\lambda_i$  to finding good rational approximations to  $\alpha$  with both numerator and denominator being *evenly divisible* (or *strongly even divisible*): We call a sequence  $m_n$  of numbers evenly divisible if there exist divisors  $d_n \mid m_n$  such that  $\min(d_n, m_n/d_n) \gg_\varepsilon m_n^{1/2-\varepsilon}$ , for all  $\varepsilon > 0$ . We call the sequence strongly evenly divisible if  $\min(d_n, m_n/d_n) \gg m_n^{1/2}$ . Then by [1, Lemma 3.1], we have:

**Lemma 1.3.** *If  $\alpha > 0$  has infinitely many rational approximations  $p_n/q_n$  satisfying*

$$(1.2) \quad |q_n \alpha - p_n| \ll \frac{1}{q_n}$$

*with the sequences  $\{p_n\}, \{q_n\}$  being evenly divisible (resp. strongly evenly divisible), then  $\delta_{\min}^{(\alpha)}(N) \ll N^{-1+\varepsilon}$  for all  $\varepsilon > 0$  (resp.  $\delta_{\min}^{(\alpha)}(N) \ll N^{-1}$ ) infinitely often.*

*If in addition  $q_n \gg q_{n+1}$  for all  $n$ , then these inequalities hold for all  $N$ .*

The authors of [1] were able to find evenly divisible approximations satisfying (1.2) for quadratic irrationalities of the form (1.1). However, the condition (1.2) can be significantly weakened; namely, if we replace it with the condition

$$(1.3) \quad |q_n \alpha - p_n| \ll \frac{1}{q_n^{1-\varepsilon}}, \text{ for all } \varepsilon > 0$$

The conclusion of Lemma 1.3 that  $\delta_{\min}^{(\alpha)}(N) \ll N^{-1+\varepsilon}$  for all  $\varepsilon > 0$  remains valid. This is essentially the content of [1, Lemma 3.2], and the form of the lemma that will be more useful to us.

---

<sup>1</sup>Given any  $\alpha \in K$ , this condition can be easily checked, as it is equivalent to  $\alpha$  satisfying  $N(\alpha) \in (\mathbb{Q}^\times)^2$  and  $\text{tr}(\alpha) > 0$ . We do not use this equivalence in this paper, and leave its proof as a simple exercise to the reader.

## 2. DIVISIBILITY PROPERTIES

In this section we establish some divisibility properties for traces and twisted traces of powers of a quadratic algebraic integer. These play similar roles to the Chebyshev polynomials featured in [1], and in fact generalize them.

**2.1. Divisibility properties of traces.** Let  $\omega \in \mathcal{O}_K$  be an algebraic integer with  $\text{tr } \omega \neq 0$ . Note that for any odd positive integer  $\ell$ ,  $\omega^\ell \in \mathcal{O}_K$ , hence  $\text{tr}(\omega^\ell)$  is an integer. Furthermore, it is divisible by  $\text{tr}(\omega)$ ; this follows from the fact that  $\frac{x^\ell + y^\ell}{x + y}$  is a symmetric polynomial in  $\mathbb{Z}[x, y]$ , and substituting  $\omega, \bar{\omega}$  for  $x, y$ . More generally, for any positive integer  $L$  which is odd and square-free, we have the decomposition

$$(2.1) \quad \text{tr}(\omega^L) = \Phi_L(\omega) \Psi_L(\omega),$$

where  $\Phi_L(\omega), \Psi_L(\omega) \in \mathbb{Z}$  are both integers, defined by

$$(2.2) \quad \Phi_L(\omega) = \prod_{\ell|L} \text{tr}(\omega^{L/\ell})^{\mu(\ell)}, \quad \Psi_L(\omega) = \prod_{\ell|L; \ell > 1} \text{tr}(\omega^{L/\ell})^{-\mu(\ell)}.$$

Indeed, both  $\Phi_L, \Psi_L$  are symmetric polynomials in  $\mathbb{Z}[\omega, \bar{\omega}]^2$ , hence their values must be in  $\mathbb{Z}$  for  $\omega \in \mathcal{O}_K$ .

In order to estimate the sizes of the factors  $\Phi_L, \Psi_L$ , assume further that  $|\bar{\omega}/\omega| \leq 1/2$ . Note that

$$e^{-2|x|} \leq 1 - |x| \leq |1 + x| \leq 1 + |x| \leq e^{2|x|}$$

for all  $|x| \leq 1/2$ , and therefore for any  $m \geq 1$ ,

$$|\text{tr}(\omega^m)| = |\omega^m + \bar{\omega}^m| = |\omega^m| \left| 1 + \left( \frac{\bar{\omega}}{\omega} \right)^m \right| \in |\omega^m| \cdot [e^{-2^{1-m}}, e^{2^{1-m}}],$$

hence,

$$(2.3) \quad |\text{tr}(\omega^L)| \asymp |\omega|^L, \quad |\Phi_L(\omega)| \asymp |\omega|^{\phi(L)}, \quad |\Psi_L(\omega)| \asymp |\omega|^{L-\phi(L)},$$

where the implied constants are all between  $e^{-2}$  and  $e^2$ . It immediately follows that

$$(2.4) \quad |\Phi_L(\omega)| \asymp |\text{tr}(\omega^L)|^{\frac{\phi(L)}{L}}, \quad |\Psi_L(\omega)| \asymp |\text{tr}(\omega^L)|^{1-\frac{\phi(L)}{L}}.$$

**2.2. Divisibility properties of twisted traces.** For  $\omega \in K$ , we define the *twisted trace* of  $\omega$  as

$$(2.5) \quad \tilde{\text{tr}}(\omega) := \text{tr}(\sqrt{D} \cdot \omega) = (\omega - \bar{\omega})\sqrt{D} \in \mathbb{Q}.$$

Suppose  $\omega \in \mathcal{O}_K$ , and  $\tilde{\text{tr}}(\omega) \neq 0$  (i.e.  $\omega \notin \mathbb{Q}$ ). Then for all positive integers  $\ell$ ,  $\tilde{\text{tr}}(\omega^\ell)$  is an integer and divisible by  $\text{tr}(\omega)$ , since  $\frac{x^\ell - y^\ell}{x - y}$  is a symmetric

---

<sup>2</sup> Specifically,  $\Phi_L$  is the homogenized reflected  $L$ -th cyclotomic polynomial, and  $\Psi_L$  is the corresponding cofactor.

polynomial in  $\mathbb{Z}[x, y]$  and  $\sqrt{D}\omega^\ell \in \mathcal{O}_K$ . As above, this generalizes to any square-free  $L \in \mathbb{N}$ , by

$$(2.6) \quad \tilde{\text{tr}}(\omega^L) = \tilde{\Phi}_L(\omega)\tilde{\Psi}_L(\omega),$$

where

$$(2.7) \quad \tilde{\Phi}_L(\omega) = \prod_{\ell|L} \tilde{\text{tr}}(\omega^{L/\ell})^{\mu(\ell)}, \quad \tilde{\Psi}_L(\omega) = \prod_{\ell|L; \ell>1} \tilde{\text{tr}}(\omega^{L/\ell})^{-\mu(\ell)}.$$

are again integers, and whenever  $|\bar{\omega}/\omega| < 1/2$  and  $L > 1$ , we have

$$(2.8) \quad |\tilde{\text{tr}}(\omega^L)| \asymp \sqrt{D}|\omega|^L, \quad |\tilde{\Phi}_L(\omega)| \asymp |\omega|^{\phi(L)}, \quad |\tilde{\Psi}_L(\omega)| \asymp \sqrt{D}|\omega|^{L-\phi(L)},$$

and consequentially

$$(2.9) \quad |\tilde{\Phi}_L(\omega)| \asymp D^{-\frac{\phi(L)}{2L}} |\tilde{\text{tr}}(\omega^L)|^{\frac{\phi(L)}{L}}, \quad |\tilde{\Psi}_L(\omega)| \asymp D^{\frac{\phi(L)}{2L}} |\tilde{\text{tr}}(\omega^L)|^{1-\frac{\phi(L)}{L}}.$$

The major difference between the divisibility properties for  $\text{tr}(\omega^L)$  and  $\tilde{\text{tr}}(\omega^L)$  is that in the second case  $L$  may be even – and specifically, we may choose  $L = 2$ , which is the only number with  $\phi(L)/L = 1/2$ , capable of generating strongly evenly divisible sequences, rather than just evenly divisible.

### 3. PROOF OF THEOREM 1.2

We first note that we may assume  $\alpha \in \mathcal{O}_K$ . Indeed, for any  $\alpha \in K$  there is some denominator  $A \in \mathbb{N}$  such that  $A \cdot \alpha \in \mathcal{O}_K$ . If we are then able to find (strongly) evenly divisible  $p_n, q_n$  such that  $A \cdot \alpha$  is well-approximated by  $p_n/q_n$ , then the sequence  $A \cdot q_n$  is also (strongly) evenly divisible and  $\alpha$  is well-approximated by  $p_n/(A \cdot q_n)$ .

Let  $\zeta \in \mathcal{O}_K$  be an algebraic integer with  $N(\zeta) = 1$  and  $|\zeta| > 1 > |\bar{\zeta}|$ . Such  $\zeta$  may constructed e.g. as  $\zeta = x + y\sqrt{D}$ , where  $(x, y) \in \mathbb{N}^2$  is a non-trivial integral solution to the Pell equation  $x^2 - Dy^2 = 1$ .<sup>3</sup> Let  $\varepsilon > 0$  be arbitrarily small.

**3.1. A symmetric construction.** Let  $\{\ell_i\}_{i=1}^\infty, \{\ell'_i\}_{i=1}^\infty$  be two disjoint subsequences of the odd primes, with

$$\prod_{i=1}^\infty \left(1 - \frac{1}{\ell_i}\right) = \prod_{i=1}^\infty \left(1 - \frac{1}{\ell'_i}\right) = \frac{1}{2},$$

the existence of which easily follows from  $\prod_p (1 - \frac{1}{p}) = 0$ . For a fixed  $t \in \mathbb{N}$  write

$$L = \prod_{i=1}^t \ell_i, \quad L' = \prod_{i=1}^t \ell'_i$$

and suppose  $t = t(\varepsilon)$  is sufficiently large so that  $\frac{\phi(L)}{L}, \frac{\phi(L')}{L'} \in (\frac{1}{2}, \frac{1}{2} + \varepsilon)$ .

---

<sup>3</sup>We may choose  $\zeta$  with  $N(\zeta) = -1$  instead, in fields where such integers exist. Note that the number  $\frac{x + \sqrt{x^2 + 4\epsilon}}{2}$  appearing in (1.1) is always an example of an appropriate  $\zeta$  for the appropriate field.

Let  $M = M(\varepsilon) \in \mathbb{N}$  be the smallest positive integer such that  $M + 1$  is divisible by  $L$  and  $M$  is divisible by  $L'$ , and write  $M + 1 = m_1 L$ ,  $M = m_2 L'$ . Define  $N = n L L'$ , where  $n = n(\varepsilon)$  is any sufficiently large number such that the following inequalities are satisfied:

$$(3.1) \quad |\zeta/\bar{\zeta}|^{nL'} > 2|\bar{\alpha}/\alpha|^{m_1}, \quad |\zeta/\bar{\zeta}|^{nL} > 2|\bar{\alpha}/\alpha|^{m_2},$$

$$(3.2) \quad |\zeta|^{\varepsilon N} > 4|\alpha - \bar{\alpha}| |N(\alpha)|^M |\alpha|^{-\varepsilon M}.$$

Such  $n$  clearly exist, as  $\alpha$  is fixed,  $m_1$ ,  $m_2$ ,  $L$  and  $L'$  depend only on  $\varepsilon$ , and  $|\zeta/\bar{\zeta}| > |\zeta| > 1$ , by the definition of  $\zeta$ . Finally, we define

$$(3.3) \quad P = P(\varepsilon) = \text{tr}(\alpha^{M+1} \zeta^N) = \text{tr}((\alpha^{m_1} \zeta^{nL'})^L),$$

$$(3.4) \quad Q = Q(\varepsilon) = \text{tr}(\alpha^M \zeta^N) = \text{tr}((\alpha^{m_2} \zeta^{nL})^{L'}).$$

We first show that  $P/Q$  is a good approximation to  $\alpha$ . Note that (3.1) implies  $|Q|$  is approximately  $|\alpha^M \zeta^N|$ , or, more explicitly,

$$(3.5) \quad |Q|/|\alpha^M \zeta^N| \in [1 - 2^{-L'}, 1 + 2^{-L'}] \subset [1/2, 2].$$

It then follows that

$$|\alpha Q - P| = |\alpha - \bar{\alpha}| |\bar{\alpha}^M \bar{\zeta}^N| = \frac{|\alpha - \bar{\alpha}| |N(\alpha)|^M |N(\zeta)|^N}{|\alpha^M \zeta^N|} \leq \frac{C(\alpha, \varepsilon)}{|Q|} \leq \frac{1}{|Q|^{1-\varepsilon}},$$

where  $C(\alpha, \varepsilon) = 2|\alpha - \bar{\alpha}| |N(\alpha)|^{M(\varepsilon)}$  depends only on  $\alpha$  and  $\varepsilon$ . The two final steps then follow immediately from (3.5), (3.2) and  $|N(\zeta)| = 1$ .

Next, we show that  $P$  and  $Q$  are evenly divisible. By their definitions, the inequalities (3.1) and section 2.1, it is evident that the factorization  $P = \Phi_L(\alpha^{m_1} \zeta^{nL'}) \Psi_L(\alpha^{m_1} \zeta^{nL'})$  satisfies

$$\min(|\Phi_L(\alpha^{m_1} \zeta^{nL'})|, |\Psi_L(\alpha^{m_1} \zeta^{nL'})|) \asymp \min(|P|^{\frac{\phi(L)}{L}}, |P|^{1-\frac{\phi(L)}{L}}) \gg |P|^{1/2-\varepsilon},$$

and similarly  $Q = \Phi_{L'}(\alpha^{m_2} \zeta^{nL}) \Psi_{L'}(\alpha^{m_2} \zeta^{nL})$  with

$$\min(|\Phi_{L'}(\alpha^{m_2} \zeta^{nL})|, |\Psi_{L'}(\alpha^{m_2} \zeta^{nL})|) \gg |Q|^{1/2-\varepsilon}.$$

We conclude that for any sequence  $\varepsilon_k \searrow 0$ , the sequences  $P(\varepsilon_k), Q(\varepsilon_k)$  are evenly divisible and provide good approximations to  $\alpha$ , which allows us to conclude via Lemma 1.3.

**3.2. Alternative constructions.** Instead of defining  $P$  and  $Q$  via traces of integral powers, we may use twisted traces instead, and define

$$P = \widetilde{\text{tr}}((\alpha^{m_1} \zeta^{nL'})^L), \quad Q = \widetilde{\text{tr}}((\alpha^{m_2} \zeta^{nL})^{L'}).$$

Repeating the same computations for the same values of  $L, L'$  would lead to the same estimates as in the first construction. However, since we are now using skew-traces, we may define  $L$  (resp.  $L'$ ) to be equal to 2 instead of a product of odd primes, which will then correspond to  $P$  (resp.  $Q$ ) having two factors of size  $\gg_D |P|^{1/2}$ , i.e. the sequence  $P(\varepsilon_k)$  (resp.  $Q(\varepsilon_k)$ ) will be strongly evenly divisible. However, we cannot in general do this for both  $P$

and  $Q$  simultaneously, since it is impossible for both  $M + 1$  and  $M$  to be even.

Suppose now that  $\alpha \in \mathbb{Q}_{>0} \cdot (K^\times)^2$ . By multiplying by a proper natural denominator  $A$ , we may assume that  $\alpha \in \mathcal{O}_K^2$ , i.e.  $\alpha = \beta^2$  for some  $\beta \in \mathcal{O}_K$ . Now, for all sufficiently large  $n$ , define

$$(3.6) \quad P_n := \widetilde{\text{tr}}(\alpha \zeta^{2n}) = \widetilde{\text{tr}}((\beta \zeta^n)^2) = \widetilde{\text{tr}}(\beta \zeta^n) \cdot \text{tr}(\beta \zeta^n),$$

$$(3.7) \quad Q_n := \widetilde{\text{tr}}(\zeta^{2n}) = \widetilde{\text{tr}}((\zeta^n)^2) = \widetilde{\text{tr}}(\zeta^n) \cdot \text{tr}(\zeta^n).$$

The sequences  $P_n$  and  $Q_n$  are therefore both strongly evenly divisible, since for large  $n$  we have

$$(3.8) \quad |\widetilde{\text{tr}}(\beta \zeta^n)| \asymp \sqrt{D} |\beta \zeta^n| \asymp \sqrt{D} |\text{tr}(\beta \zeta^n)|,$$

$$(3.9) \quad |\widetilde{\text{tr}}(\zeta^n)| \asymp \sqrt{D} |\zeta^n| \asymp \sqrt{D} |\text{tr}(\zeta^n)|.$$

Furthermore, their ratios approximate  $\alpha$  well, as

$$|Q_n \alpha - P_n| = |\alpha - \bar{\alpha}| |\bar{\zeta}|^{2n} = \frac{|\alpha - \bar{\alpha}|}{|\zeta|^{2n}} \asymp \frac{|\alpha - \bar{\alpha}| \sqrt{D}}{|Q_n|},$$

and the denominators grow geometrically –  $|Q_n| \asymp |\zeta^{-2}| |Q_{n+1}| \gg |Q_{n+1}|$ .

Therefore for such  $\alpha$ , we have  $\delta_{\min}^{(\alpha)}(N) \ll N^{-1}$  for all large  $N$ , via Lemma 1.3.

**3.3. Comparison to previous results.** For completeness, we present the constructions used in the original proof of Theorem 1.1, in terms of our notation. As mentioned above, the term  $\frac{x + \sqrt{x^2 + 4\epsilon}}{2}$  is simply our  $\zeta$ . The optional term  $\sqrt{x^2 + 4\epsilon}$  belongs to  $\sqrt{D} \cdot \mathbb{Q}^\times$ . Therefore the numbers in  $K$  covered by Theorem 1.1 are those of the form

$$\alpha(\zeta; r, a, b) = r \cdot \zeta^a \cdot \sqrt{D}^b$$

where  $\zeta \in \mathcal{O}_K^\times$ ,  $r \in \mathbb{Q}^\times$ ,  $a \in \mathbb{Z}$  and  $b = 0, 1$ . The term  $r = c/d$  is dealt with by multiplying all numerators by  $c$  and denominators by  $d$ , and we reduce to the case  $r = 1$  (and thus  $\alpha \in \mathcal{O}_K$ ). These  $\alpha$  have the special property that the sequences of traces  $\text{tr}(\alpha \zeta^N) = \text{tr}(\sqrt{D}^b \zeta^{N+a})$ , and  $\text{tr}(\zeta^N)$  (as well as  $\widetilde{\text{tr}}(\alpha \zeta^N)$  and  $\widetilde{\text{tr}}(\zeta^N)$ ) can have useful divisibility properties, for good choices of  $N$ , and their ratios provide good approximations to  $\alpha$ . For general  $\alpha$ , we can generalize this method and generate good approximations to  $\alpha$  as ratios of the sequences of  $\text{tr}(\alpha \omega \zeta^N)$  and  $\text{tr}(\omega \zeta^N)$ , for any  $\omega \in \mathcal{O}_K$ . The quality of the approximation is slightly worse: the upper bound on the error grows by the norm of  $\omega$ , but this is not a problem if  $\omega$  is fixed and  $N$  is large. In order to have good divisibility properties for these sequences we want  $\alpha \omega \zeta^N$  and  $\omega \zeta^N$  to be  $L$ -th and  $L'$ -th powers, respectively. We achieve this by setting  $\omega = \alpha^M$  and choosing appropriate values of  $M$  and  $N$  – but other choices for  $\omega$  and  $N$  might also be possible.

## REFERENCES

- [1] BLOMER, V., BOURGAIN, J., RADZIWIŁŁ, M., AND RUDNICK, Z. Small gaps in the spectrum of the rectangular billiard. arXiv:1604.02413v3. To appear in Ann. Sci. École Norm. S..

RAYMOND AND BEVERLY SACKLER SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV  
UNIVERSITY, TEL AVIV 69978, ISRAEL